



ELSEVIER

Nuclear Instruments and Methods in Physics Research B 182 (2001) 193–199

---

---

**NIM B**  
Beam Interactions  
with Materials & Atoms

---

---

www.elsevier.com/locate/nimb

# Reactive ion scattering from pure and mixed HCl, NH<sub>3</sub> and D<sub>2</sub>O surfaces

Seong-Chan Park <sup>a</sup>, Kye-Won Maeng <sup>a</sup>, T. Pradeep <sup>b,\*</sup>, H. Kang <sup>a,\*</sup><sup>a</sup> Department of Chemistry and Center for Ion-Surface Reaction, Pohang University of Science and Technology,  
Pohang 790-784, Republic of Korea<sup>b</sup> Department of Chemistry and Regional Sophisticated Instrumentation Centre, Indian Institute of Technology,  
Madras 600 036, India

---

## Abstract

We have sampled the various species present on the condensed molecular layers of HCl, NH<sub>3</sub>, D<sub>2</sub>O and their mixtures by Cs<sup>+</sup> reactive ion scattering (RIS) and low-energy sputtering. HCl and NH<sub>3</sub> exist in molecular states on pure molecular films, but the ammonium ion (NH<sub>4</sub><sup>+</sup>) is readily formed when HCl and NH<sub>3</sub> are co-deposited, by proton transfer from HCl to NH<sub>3</sub>. Water is unnecessary for the proton transfer, although enhances it. In the presence of water, HCl ionizes first to form the hydronium ion (H<sub>3</sub>O<sup>+</sup>), which subsequently transfers the proton to NH<sub>3</sub> to form NH<sub>4</sub><sup>+</sup>. This work demonstrates the capability of RIS for probing chemical species on condensed molecular surfaces and monitoring their reactions. © 2001 Elsevier Science B.V. All rights reserved.

*Keywords:* Low-energy ion scattering; Ice; Ammonia; Hydrogen chloride; Cesium; Ruthenium

*PACS:* 34.50.D; 61.72.S; 82.65.J; 92.60.S

---

## 1. Introduction

There is an increasing interest for studying reactions occurring on cold ice surfaces. Such interest arises from the recent discovery which suggests that the surfaces of cold ice particles may play impor-

tant roles in the chemistry of the Earth's atmosphere [1] and formation of organic molecules in the interstellar space [2]. Current understanding of reactions on ice, however, is very limited, compared to that in aqueous media which has been pursued since the beginning of chemistry. For instance, a simple process like ionization of HCl on cold ice surface is not clearly understood to the molecular details, despite recent active investigations using various experimental and theoretical approaches [3–7]. The molecular and ionized states of HCl on ice have recently been identified by Cs<sup>+</sup> reactive ion scattering (RIS) technique, and their relative portions have been quantified [7]. RIS is a


---

\* Corresponding author. Present address: Department of Chemistry, Seoul National University, Kwanak-ku, Shinrim-dong, Seoul 151-742, South Korea. Tel.: +82-2-875-74-71; fax: +82-2-889-57-19.

E-mail addresses: pradeep@iitm.ac.in (T. Pradeep), surf-ion@snu.ac.kr (H. Kang).

sensitive tool to monitor species at surfaces [8–10]; the surface layer investigated is extremely thin, corresponding to depths of only one to two atomic layers in the energies employed (5–100 eV). In this study, we use the RIS technique to examine the chemical states of HCl, NH<sub>3</sub>, D<sub>2</sub>O, and their mixtures frozen onto a cold substrate. Proton transfer of HCl to NH<sub>3</sub> in the frozen layers is examined in the presence and absence of water molecules.

The HCl–NH<sub>3</sub> system is a model case to study a proton transfer phenomenon and has long attracted attention. The reaction



in aqueous solutions is instantaneous with an equilibrium constant ( $K_{\text{eq}}$ ) of  $1.7 \times 10^9$  at 298 K, and is one of the textbook examples of acid–base chemistry. The role of solvent molecules in the proton transfer of HCl to NH<sub>3</sub> has been a question of great academic importance. Early calculations [11–13] lend support to the ionic character of the interaction in the gas phase. The Van der Waals complex, NH<sub>3</sub> ··· HCl has been detected in a microwave spectroscopy experiment [14,15], and several computational investigations supported the existence of such a molecular system [16–21]. The situation, however, becomes drastically different once water molecules are added to this complex. While it is still a Van der Waals molecule with a flat potential energy pathway for proton transfer with one water molecule, two and three molecules make it an ion pair [22]. Addition of water molecules increases the stability of the ion pair relative to the hydrogen-bonded system. Nevertheless, it is well known that particles of NH<sub>4</sub>Cl are formed when dry vapors of NH<sub>3</sub> and HCl are mixed, which has been explained in terms of the heterogeneous chemistry on liquid or solid particles pre-existing in the system. In a recent calculation, Tao [23] showed that while proton transfer is not possible in a single NH<sub>3</sub>–HCl system, in clusters containing two and four hetero-dimer units, complete proton transfer occurs by maximizing electrostatic interactions. This homogeneous nucleation may have implications to the processes in aerosols and clouds [24].

## 2. Experiment

In the RIS experiment, a Cs<sup>+</sup> ion beam produced from a surface ionization source (Kimball Physics) was scattered from a sample being examined at the desired kinetic energy (5–100 eV). The Cs<sup>+</sup> current density at the target was 1–5 nA/cm<sup>2</sup>. The positive ions emitted from the sample surface were mass-analyzed by a quadrupole mass spectrometer (QMS) with its ionizer filament turned off. The emitted ions are composed of reflected primaries, RIS products which are association products of Cs<sup>+</sup> with the neutrals at the surface, and pre-existing ions sputtered by the low-energy Cs<sup>+</sup> impact. Each ion mass spectrum consisted of at least 10 scans which took about 15 s to acquire. The substrate was an Ru single crystal with a (001) face, and its temperature could be varied in the range 90–1300 K on a variable-temperature sample stage. The beam incidence and the detector angles were both 67.5° with respect to the surface normal. The chamber has separate dosing facilities for gases. Frozen water films were prepared by exposing D<sub>2</sub>O vapor on an Ru(001) substrate kept at 120 K. At this temperature, the ice layer will have a non-porous, amorphous structure [25]. Frozen NH<sub>3</sub> and HCl layers were made by condensing the corresponding gases at 100 K, unless specified otherwise. The chamber has a base vacuum of  $3 \times 10^{-10}$  Torr and is equipped with an Auger spectrometer to monitor surface cleanliness and surface composition. Detailed description for the UHV chamber can be found elsewhere [9,10]. The QMS for scattered ion analysis could also be operated in the residual gas analysis mode for thermal desorption spectrometry (TDS).

## 3. Results

In Fig. 1, we show the positive ion mass spectra of the scattering experiment performed on the frozen (a) NH<sub>3</sub> and (b) HCl layers deposited on a Ru(001) surface. The Cs<sup>+</sup> collision energy was 30 eV. Spectrum (a) shows a strong, elastic peak of Cs<sup>+</sup> at  $m/z = 133$  amu/charge, together with RIS peaks at  $m/z = 150$  (CsNH<sub>3</sub><sup>+</sup>) and 167 amu/charge




Fig. 1. Scattered ion mass spectra upon the collision of Cs<sup>+</sup> on pure NH<sub>3</sub> (a) and pure HCl multilayers (b), and NH<sub>3</sub>-multilayer exposed to 0.5 L HCl gas (c). Each layer was prepared by condensing the corresponding gases on an Ru(001) surface at 100 K through separate tube dosers. The multilayer formation was checked by TDS and Auger spectroscopy. The species observed are labeled. Cs<sup>+</sup> impact energy was 30 eV.

(CsN<sub>2</sub>H<sub>6</sub><sup>+</sup>). The CsNH<sub>3</sub><sup>+</sup> peak is formed by pickup of an NH<sub>3</sub> molecule from the surface, and CsN<sub>2</sub>H<sub>6</sub><sup>+</sup> by pickup of two NH<sub>3</sub> molecules. The RIS process leading to Cs<sup>+</sup>-molecule clusters at condensed molecular surfaces has been described in detail in a previous paper on water-ice surfaces [26]. In spectrum (a) no secondary ions due to low-energy sputtering were seen in the region below 133 amu/charge. The appearance of the CsNH<sub>3</sub><sup>+</sup> peak and the absence of low-energy sputtered ions




Fig. 2. RIS mass spectra obtained on D<sub>2</sub>O-ice surface exposed to 0.5 L of HCl (a) and 2 L of NH<sub>3</sub> (b) at 140 K. The D<sub>2</sub>O-ice layer was 3–4 bilayers (BL) thick, prepared on an Ru (001) surface at 120 K through D<sub>2</sub>O backfilling. The exposures for HCl and NH<sub>3</sub> were controlled to give approximately the same concentrations of these species on the surfaces.

indicate that ammonia exists as a neutral molecule on the frozen surface. Fig. 2(b) presents an RIS spectrum from a frozen HCL layer. The RIS peak at m/z = 169 (CsHCl<sup>+</sup>) represents pickup of molecular HCl on the surface. The peak at m/z = 161 (CsH<sub>2</sub>O<sup>+</sup>) is due to residual water molecules inside the chamber that were deposited on the surface. In the low-mass region, secondary ion signals were observed at m/z = 19 (H<sub>3</sub>O<sup>+</sup>), indicating the presence of protonated water species on the surface produced by HCl ionization aided by deposited residual water species [7]. It may be noted that H<sub>3</sub>O<sup>+</sup> ions are not emitted from a pure H<sub>2</sub>O-ice surface by low-energy sputtering [26], thus excluding the possibility that the H<sub>3</sub>O<sup>+</sup> signal is produced by impact-induced ionization of water molecules.

In Fig. 1(c), we exposed 0.5 L of HCl onto an NH<sub>3</sub>-predeposited surface. The NH<sub>3</sub> film was deliberately kept thick so that there is no effect

caused by the interaction of HCl and the underlying Ru substrate. The first significant aspect to be noticed is the large  $\text{NH}_4^+$  secondary ion intensity (note the reduced scale) which is quite unusual in such spectra. In addition, we also see the ammoniated ammonium ion,  $\text{N}_2\text{H}_7^+$ . The RIS products observed are  $\text{CsNH}_3^+$ ,  $\text{CsN}_2\text{H}_6^+$ ,  $\text{CsHCl}^+$ , and  $\text{CsNH}_4\text{Cl}^+$ . The  $\text{NH}_4^+$  species is solely due to the ejection of preformed ions as revealed by threshold measurements; while  $\text{NH}_4^+$  had threshold energy of 17 eV on the HCl/ $\text{NH}_3$  surface, it was above 60 eV on a pure  $\text{NH}_3$  surface. The strong  $\text{NH}_4^+$  intensity, far greater than  $\text{H}_3\text{O}^+$  in Fig. 1(b), reveals that  $\text{NH}_4^+$  is formed by proton transfer from HCl to  $\text{NH}_3$  on the frozen  $\text{NH}_3$  layer at 100 K. Apparently, water is unnecessary to assist the proton transfer.

We examined the chemical states of HCl and  $\text{NH}_3$  deposited on water–ice films. Fig. 2 compares the RIS spectra obtained when (a) HCl and (b)  $\text{NH}_3$  gases were deposited in a submonolayer coverage on  $\text{D}_2\text{O}$ –ice films at 140 K. As mentioned before, in both spectra, multilayers of  $\text{D}_2\text{O}$  were deposited to avoid any possible substrate effect in the measurement. In Fig. 2(a), the peak at  $m/z = 153$  represents  $\text{CsD}_2\text{O}^+$ , and its nearby peaks are H/D exchange products.  $\text{CsHCl}^+$  peak is absent due to complete ionization of HCl on ice at 140 K [7]. The secondary ion peaks include  $\text{HD}_2\text{O}^+$ ,  $\text{HD}_4\text{O}_2^+$ , and their H/D-exchanged families. These spectral features confirm ionic dissociation of HCl reported previously [7]. In Fig. 1(b), where  $\text{NH}_3$  is deposited on ice, no protonated species appear in the low-mass region. The RIS products include  $\text{CsNH}_3^+$ ,  $\text{CsD}_2\text{O}^+$ , and  $\text{Cs}(\text{NH}_3)_x(\text{D}_2\text{O})_y^+$  ( $x, y = 1, 2$ ). The H/D exchange does not occur for these species. These results confirm that  $\text{NH}_3$  remains unionized on ice.

Sticking probability of  $\text{NH}_3$  and HCl gases at an ice surface is not well known at the temperatures employed in this study, and thus it is desirable to estimate the actual coverage of the species deposited on ice surfaces. We obtained  $\text{NH}_3$ -deposition curves by measuring  $\text{CsNH}_3^+$  intensity as a function of  $\text{NH}_3$  exposure on ice. Fig. 3 exemplifies the deposition curve, showing the variation of the intensity ratio of  $\text{CsNH}_3^+$  to  $\text{CsD}_2\text{O}^+$  against  $\text{NH}_3$  exposure at 100 K. The plot is approximately




Fig. 3. Plot of the intensity ratio of  $\text{CsNH}_3^+$  to  $\text{CsD}_2\text{O}^+$  against  $\text{NH}_3$  exposure.  $\text{NH}_3$  gas had a partial pressure of  $1 \times 10^{-8}$  Torr during the exposure. Fresh surfaces were prepared for each data point to remove the Cs contamination effect from the beam.

linear for  $\text{NH}_3$  exposure up to 200 s at a partial pressure of  $1 \times 10^{-8}$  Torr, indicating a constant sticking probability of  $\text{NH}_3$  over this span. Since the sampling depth of RIS is 1–2 monolayers of ice at the present beam energy, the linearity of the curve also indicates that  $\text{NH}_3$  has a fractional coverage on the surface.

We examined the reaction between  $\text{NH}_3$  and HCl occurring on  $\text{D}_2\text{O}$ –ice. In Fig. 4(a), the surface was prepared by exposing a multilayer film of  $\text{D}_2\text{O}$ –ice to 0.5 L of HCl gas at 100 K, and then to 0.3 L of  $\text{NH}_3$  gas. Secondary ion peaks include both ammonium and hydronium species. Apparently, hydronium ions are formed by ionic dissociation of HCl, as mentioned before, which then react with  $\text{NH}_3$  to generate ammonium ions by proton transfer described in Reaction (1).



RIS peaks include  $\text{CsD}_2\text{O}^+$  and  $\text{CsNH}_3^+$ , indicating the presence of  $\text{D}_2\text{O}$  and  $\text{NH}_3$  molecules on the surface as well. Thus, all four species of Reaction




Fig. 4. Result of RIS measurement for the proton transfer reaction,  $\text{H}_3\text{O}^+ + \text{NH}_3 \leftrightarrow \text{H}_2\text{O} + \text{NH}_4^+$ . (a) Hydronium ions were generated on a  $\text{D}_2\text{O}$ -ice surface by 0.5 L exposure of HCl gas, which then were reacted with  $\text{NH}_3$  gas for 0.3 L exposure. (b) Ammonium ions were first generated by exposing 0.2 L HCl to an  $\text{NH}_3$  multilayer, and then 0.5 L of  $\text{D}_2\text{O}$  was added. The underlying multilayer films were grown at 120 K, and the acid-base titration experiments were carried out at 100 K.

(1) are present on the surface as identified by the spectrum. They all exhibit substantial intensities, indicating that Reaction (1) is incomplete.

The reverse process of Reaction (1) was also investigated, i.e., the proton transfer from  $\text{NH}_4^+$  to  $\text{D}_2\text{O}$ . In Fig. 4(b), an  $\text{NH}_3$  multilayer was prepared without water on Ru(001) and then exposed to HCl to produce  $\text{NH}_4^+$  ions, as was done in Fig. 1(c).  $\text{D}_2\text{O}$  was then exposed to this surface. The spectrum shows an intense  $\text{NH}_4^+$  peak, but an extremely small intensity for  $\text{HD}_2\text{O}^+$ .  $\text{CsNH}_3^+$  and  $\text{CsD}_2\text{O}^+$  peaks are also seen. The  $\text{CsNH}_3^+$  peak must come from the underlying  $\text{NH}_3$  layer, rather than from the reverse reaction.  $\text{CsHDO}^+$  must be due to H/D exchange facilitated by HCl ionization in the presence of  $\text{D}_2\text{O}$ . The strong  $\text{NH}_4^+$  signal and the almost absent  $\text{HD}_2\text{O}^+$  signal indicate that

the reverse proton transfer does not take place to any significant degree.

#### 4. Discussion

The spectral features described in Section 3 reveal the chemical states of  $\text{NH}_3$ , HCl, and  $\text{D}_2\text{O}$  on the frozen layers.  $\text{NH}_3$  exists as a neutral molecule without protonation both in the presence and absence of water molecules (Figs. 1(a) and 2(b)). This observation is well expected since  $\text{NH}_3$  is a weak base in the aqueous phase as well. HCl is believed to be in a molecular form on a frozen HCl surface, and the small intensity of  $\text{H}_3\text{O}^+$  signal in Fig. 1(b) is attributed to residual water vapor. HCl efficiently ionizes on  $\text{D}_2\text{O}$ -ice at temperature above 100 K, with the ionization degree varying with ice temperature [7]. When  $\text{NH}_3$  and HCl are co-deposited to a frozen film, proton transfer reaction occurs from HCl to  $\text{NH}_3$  to produce  $\text{NH}_4^+$  (Fig. 1(c)). Since the  $\text{NH}_4^+$  signal in Fig. 1(c) is far larger than the  $\text{H}_3\text{O}^+$  signal in Fig. 1(b), it is clear the proton transfer readily occurs even in the absence of water molecules. Neighboring  $\text{NH}_3$  molecules can assist the reaction by clustering and maximizing HCl- $\text{NH}_3$  electrostatic interactions. The enhanced proton transfer by solvating  $\text{NH}_3$  molecules was predicted by theoretical calculation [23]. On a  $\text{D}_2\text{O}$ -ice surface (Fig. 4(a)), ammonium ions are also efficiently formed upon co-deposition of  $\text{NH}_3$  and HCl. In this case, the ammonium species are created in a large part also by the proton transfer from hydronium ion to ammonia (Reaction (1)), the former generated by HCl ionization on  $\text{D}_2\text{O}$ -ice. When  $\text{NH}_3$  and HCl build up sufficiently high surface concentrations on the surface, the direct proton transfer between the two species may also become possible.

An interesting species observed from the  $\text{NH}_3$ -HCl mixture (Fig. 1(c)) is the  $\text{CsNH}_4\text{Cl}^+$  peak, which represents an association product of  $\text{Cs}^+$  with  $\text{NH}_4\text{Cl}$ .  $\text{NH}_4\text{Cl}$  corresponds to either  $\text{NH}_4\text{Cl}$  salt or  $\text{NH}_3 \cdots \text{HCl}$  Van der Waals complex. This ion is not seen or the intensity is very small as water is introduced at the surface (Fig. 4). In this respect, the  $\text{CsNH}_4\text{Cl}^+$  signal in Fig. 1(c) may suggest the presence of  $\text{NH}_3 \cdots \text{HCl}$  van der

Waals complex at the surface. The stability of the van der Waals complex will decrease in the presence of water, as expected from thermodynamics. It is also possible that  $\text{NH}_4\text{Cl}$  salt is formed on the frozen  $\text{NH}_3\text{--HCl}$  surface surrounded by ammonia molecules, as suggested by  $\text{NH}_4^+$  peak, which then is pick up by  $\text{Cs}^+$ . Theoretical calculations [16–23], however, predict that a Van der Waals complex will be more stable when  $\text{NH}_4\text{Cl}$  unit exists as an isolated molecule.

Fig. 4 allows us not only to identify all the components of Reaction (1), but also to measure their relative concentrations at the surfaces. We can express the quotient ( $Q$ ) of Reaction (1) by the peak intensities according to the following equation:

$$Q = \frac{[\text{H}_2\text{O}][\text{NH}_4^+]}{[\text{H}_3\text{O}^+][\text{NH}_3]} \\ = \frac{I(\text{CsH}_2\text{O}^+)I(\text{NH}_4^+)}{I(\text{H}_3\text{O}^+)I(\text{CsNH}_3^+)} \quad (2)$$

Here  $I(\text{X}^+)$  represents signal intensity of  $\text{X}^+$ . Note that the relationship of Eq. (2) is quantitative. Although the detection sensitivities are different between the protonated  $\text{HX}^+$  ions (detected as secondary ions) and the X neutrals (detected as  $\text{Cs}^+$  RIS products), the different sensitivity factors will cancel out automatically in the expression of Eq. (2) because they appear both in the numerator and denominator. The  $Q$  value thus calculated is about 20 in the  $\text{NH}_3\text{--HCl}$  titration experimental of Fig. 4(a).

The measured  $Q$  value is much smaller than the equilibrium constant of Reaction (1) in the aqueous phase ( $1.7 \times 10^9$ ). The gas-phase equilibrium constant for Reaction (1) is even larger ( $1 \times 10^{30}$ ), which is calculated from the gas-phase proton affinities,  $\text{PA}(\text{H}_2\text{O}) = 166$  kcal/mol and  $\text{PA}(\text{NH}_3) = 207$  kcal/mol. Since reaction on ice is characterized by partial solvation of reactants and products by water molecules, according to the theoretical works that investigated HCl ionization on ice [5], we may well expect that  $K_{\text{eq}}$  of the reaction on ice will be between the gas-phase and aqueous-phase values. The present observation, however, shows that this is not the case, indicating that Reaction (1) at an ice surface does not reach a thermodynamic equilibrium but is in a metastable state.

## 5. Conclusion

We demonstrated here that RIS provides a suitable means for examining the chemical states of condensed molecular surfaces and for quantitatively monitoring their relative concentration. Neutral states of HCl and  $\text{NH}_3$  were unambiguously identified from the RIS signals, and their ionized forms were detected by low-energy sputtering. Interesting feature of the experiment is that the acid–base reaction occurring right on the monolayer surface can be monitored. The simple evaluation of the quotient for the  $\text{H}_3\text{O}^+\text{--NH}_3$  reaction on ice revealed that the reaction does not reach a thermodynamic equilibrium but is in a metastable state. Such a study may be extendible to the determination of the parameters appropriate for quantitative acid–base chemistry, such as proton affinity and acidity scale of ice surfaces. A complete study of the reaction aiming to deduce these quantities, including examination of the effect of ice temperature, morphology, reactant concentration, and solvation and subsurface diffusion of reactant species, is now underway.

## Acknowledgements

The work was supported by the Creative Research Initiatives Project from the Korea Institute of Science & Technology Evaluation and Planning.

## References

- [1] M.J. Molina, T.L. Tso, L.T. Molina, F.C.Y. Wang, *Science* 238 (1987) 1253.
- [2] M.P. Bernstein et al., *Science* 283 (1999) 1135.
- [3] S.F. Banham, A.B. Horn, T.G. Koch, J.R. Sodeau, *Faraday Discuss.* 100 (1995) 321.
- [4] L. Delzeit, B. Rowland, J.P. Devlin, *J. Phys. Chem.* 97 (1993) 10, 312.
- [5] B.J. Gertner, J.T. Hynes, *Science* 271 (1996) 1563.
- [6] C.J. Pursell, M. Zaidi, A. Thompson, C. Fraser-Gaston, E. Vela, *J. Phys. Chem. A* 104 (2000) 552.
- [7] H. Kang, T.-H. Shin, S.-C. Park, I.K. Kim, S.-J. Han, *J. Am. Chem. Soc.* 122 (2000) 9842.
- [8] R.G. Cooks, T. Ast, T. Pradeep, V.H. Wysocki, *Acc. Chem. Res.* 27 (1994) 316.

- [9] M.C. Yang, C.H. Hwang, H. Kang, *J. Chem. Phys.* 107 (1997) 2611.
- [10] K.Y. Kim, T.H. Shin, S.-J. Han, H. Kang, *Phys. Rev. Lett.* 82 (1999) 1329.
- [11] R.S. Mulliken, *J. Phys. Chem.* 56 (1952) 801.
- [12] R.S. Mulliken, *Science* 157 (1967) 13.
- [13] E. Clementi, *J. Chem. Phys.* 46 (1967) 3851.
- [14] E.J. Goodwin, N.W. Howard, A.C. Legon, *Chem. Phys. Lett.* 131 (1986) 319.
- [15] N.W. Howard, A.C. Legon, *J. Chem. Phys.* 88 (1988) 4694.
- [16] Z. Latajka, S. Sakai, K. Morokuma, H. Ratajczak, *Chem. Phys. Lett.* 110 (1984) 464.
- [17] A. Brciz, A. Karpfen, H. Lischka, P. Schuster, *Chem. Phys.* 89 (1984) 337.
- [18] P.G. Jasien, W.J. Stevens, *Chem. Phys. Lett.* 130 (1986) 127.
- [19] C. Chipot, D. Rinaldi, J.-L. Rivail, *Chem. Phys. Lett.* 191 (1992) 287.
- [20] G.B. Bacskay, *Mol. Phys.* 77 (1992) 61.
- [21] G. Corongju, D. Estrin, G. Murgia, L. Paglieri, L. Pisani, G. Suzzi Valli, J.D. Watts, E. Clementi, *Int. J. Quantum Chem.* 59 (1996) 119.
- [22] M. Biczysko, Z. Latajka, *Chem. Phys. Lett.* 313 (1999) 366.
- [23] F.-M. Tao, *J. Chem. Phys.* 110 (1999) 11,121.
- [24] D.W. Oxtoby, *Acc. Chem. Res.* 31 (1998) 91.
- [25] D.E. Brown, S.M. George, C. Huang, E.K. Wong, K.B. Rider, R.S. Smith, B.D. Kay, *J. Phys. Chem.* 100 (1996) 4988.
- [26] T.H. Shin, S.-J. Han, H. Kang, *Nucl. Instr. and Meth. B* 157 (1999) 191.